



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

A retrospective study of bacterial pathogens in an equine hospital (1988–2014)

van Spijk, Julia N ; Schmitt, S ; Fürst, Anton E ; Schoster, Angelika

Abstract: Knowledge about local pathogen prevalence is important for guiding initial antimicrobial therapy. The aim of this study was to describe the longitudinal occurrence of bacterial pathogens in hospitalized horses at the University of Zurich between 1988 and 2014. Positive culture results were retrospectively analyzed. Isolates were grouped based on sample site, diagnosis, and year of isolation. The frequency of bacterial isolates over time was compared using the Chi-Square test. *Escherichia coli* (299/1,723, 17%), *Streptococcus equi* ssp. *zooepidemicus* (295/1,723, 17%) and coagulase-positive staphylococci (196/1,723, 11%) were the most common pathogens. Anaerobes were common in dental (18/66, 27%), peritoneal (6/43, 14%) and soft tissue infections (28/208, 13%). Mixed infections were common in skin and dental infections (24/55, 44% and 17/43, 40%, respectively). Frequently isolated bacteria in most organ systems included Gram-positives and Gram-negatives, supporting the need for initial broad-spectrum antimicrobial therapy until culture and susceptibility results become available.

DOI: <https://doi.org/10.17236/sat00068>

Other titles: Eine retrospektive Studie pathogener Bakterien bei hospitalisierten Pferden (1988–2014)

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-124663>

Journal Article

Published Version

Originally published at:

van Spijk, Julia N; Schmitt, S; Fürst, Anton E; Schoster, Angelika (2016). A retrospective study of bacterial pathogens in an equine hospital (1988–2014). *Schweizer Archiv für Tierheilkunde*, 158(6):423–431.

DOI: <https://doi.org/10.17236/sat00068>

A retrospective study of bacterial pathogens in an equine hospital (1988–2014)

J. N. van Spijk¹, S. Schmitt², A. E. Fürst¹, A. Schoster¹

¹Equine Department and ²Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Zurich, Switzerland

Abstract

Knowledge about local pathogen prevalence is important for guiding initial antimicrobial therapy. The aim of this study was to describe the longitudinal occurrence of bacterial pathogens in hospitalized horses at the University of Zurich between 1988 and 2014. Positive culture results were retrospectively analyzed. Isolates were grouped based on sample site, diagnosis, and year of isolation. The frequency of bacterial isolates over time was compared using the Chi-Square test. *Escherichia coli* (299/1,723, 17%), *Streptococcus equi* ssp. *zooepidemicus* (295/1,723, 17%) and coagulase-positive staphylococci (196/1,723, 11%) were the most common pathogens. Anaerobes were common in dental (18/66, 27%), peritoneal (6/43, 14%) and soft tissue infections (28/208, 13%). Mixed infections were common in skin and dental infections (24/55, 44% and 17/43, 40%, respectively). Frequently isolated bacteria in most organ systems included Gram-positives and Gram-negatives, supporting the need for initial broad-spectrum antimicrobial therapy until culture and susceptibility results become available.

Keywords: horse, hospitalized patients, bacteria, retrospective long-term analysis

Eine retrospektive Studie pathogener Bakterien bei hospitalisierten Pferden (1988–2014)

Kenntnisse über lokal vorkommende pathogene Bakterien sind wichtig, um initial eine gezielte antibiotische Therapie einleiten zu können. Das Ziel der Studie war es, die häufigsten Infektionserreger und deren zeitliche Veränderungen bei Pferden am Tierspital Zürich zwischen 1988 und 2014 zu beschreiben. Alle positiven Kulturergebnisse wurden retrospektiv analysiert. Die Isolate wurden anhand ihrer Lokalisation, Diagnose und Jahr der Isolation eingeteilt. Die Häufigkeit verschiedener Keime wurde mittels des Chi-Square Tests zwischen verschiedenen Zeitphasen verglichen. *Escherichia coli* (299/1,723, 17%), *Streptococcus equi* ssp. *zooepidemicus* (295/1,723, 17%) und koagulasepositive Staphylokokken (196/1,723, 11%) waren die insgesamt am häufigsten nachgewiesenen Bakterien. Obligat anaerobe Bakterien wurden vorwiegend bei Zahninfektionen (18/66, 27%), Peritonitis (6/43, 14%) und Weichteilinfektionen (28/208, 13%) isoliert. Mischkulturen waren besonders häufig bei Haut- und Zahninfektionen (24/55, 44% und 17/43, 40%). Zu den am häufigsten isolierten Keimen gehörten in den meisten Organsystemen sowohl Gram-positive als auch Gram-negative Erreger. Daher sollte die Wahl des Antibiotikums bis zum Vorliegen der Kulturresultate und Antibiotogramme auf ein Medikament mit breitem Wirkspektrum fallen.

Schlüsselwörter: Pferd, hospitalisierte Patienten, Bakterien, retrospektive Langzeitstudie


DOI 10.17236/sat00068

Received: 12.01.2016

Accepted: 12.03.2016

Table 1: Origin of bacterial pathogens isolated from horses in Zurich between 1988 and 2014.

| Organ system and diagnosis | Number of isolates per organ system (%) | Number of isolates per diagnosis (%) |
|-------------------------------|-----------------------------------------|--------------------------------------|
| Respiratory | 397 (23%) | |
| Lower airway infection | | 256 (64%) |
| Upper airway infection | | 141 (36%) |
| Post-procedural | 386 (22%) | |
| Post-procedural (soft tissue) | | 267 (69%) |
| Post-procedural (orthopedic) | | 106 (28%) |
| Post-procedural (sinus) | | 13 (3%) |
| Soft tissue | 208 (12%) | |
| External abscess | | 110 (53%) |
| Soft tissue wound | | 52 (25%) |
| Cellulitis | | 41 (20%) |
| Infected hematoma | | 4 (2%) |
| Myositis | | 1 (<1%) |
| Musculoskeletal | 207 (12%) | |
| Synovial infection | | 96 (46%) |
| Osteitis | | 72 (35%) |
| Hoof abscess | | 39 (19%) |
| Reproductive | 99 (6%) | |
| Genital tract infection | | 97 (98%) |
| Mastitis | | 2 (2%) |
| Urinary | 95 (6%) | |
| Urinary tract infection | | 67 (71%) |
| Omphalitis | | 27 (28%) |
| Renal abscess | | 1 (1%) |
| Skin | 87 (5%) | |
| Dental | 66 (4%) | |
| Peritoneal | 43 (3%) | |
| Sepsis | 27 (2%) | |
| Nervous system | 3 (<1%) | |
| Others | 108 (6%) | |
| Total | 1723 (100%) | |

**Figure 1:** Distribution of equine samples (n=1,281) with a single species of pathogenic bacteria and proportion of mixed infections by organ system isolated in Zurich between 1988 and 2014.

Introduction

Bacterial infections are a major cause of morbidity and mortality in horses. The identification of pathogenic bacteria in disease processes is important for understanding pathogenesis and optimizing therapy (Sellon and Long, 2013). While clinical findings often allow a presumptive diagnosis of bacterial infection, confirmation and characterization of the causative agent is achieved through bacteriological culture and susceptibility testing (Morley et al., 2005). However, culture results are often not available for up to 48 hours, and an empiric choice of antimicrobial treatment has to be made (Wilson, 2001). To guide the choice of initial antimicrobial therapy, knowledge of common pathogens in affected organ systems as well as of antimicrobial resistance of common pathogens is important (Wilson, 2001). There are studies available describing bacterial isolates from equine infections, however, only few of them are recent (Hirsh and Jang, 1987; Lavoie et al., 1991; Sweeney et al., 1991; Hawkins et al., 1993; Racklyeft and Love, 2000; Clark et al., 2008; Panchaud et al., 2010; Theelen et al., 2014). Additionally, bacterial prevalence can differ geographically; therefore, surveillance of bacterial isolates is recommended on a local level (Morley et al., 2005; Traub-Dargatz and Dargatz, 2009). There is currently little data available of common bacterial pathogens in horses in Switzerland.

In recent years, antimicrobial drug resistance has become an emerging problem in human and veterinary medicine (Weese, 2009). The acquisition of resistance against antimicrobials permits survival of particular bacterial strains, and thus changes in susceptibility profiles are able to shift the prevalence of pathogens. Therefore, surveillance of changes in prevalence is recommended (Morley et al., 2005; Traub-Dargatz and Dargatz, 2009). The objective of this study was to retrospectively describe occurrence and longitudinal development of bacterial pathogens isolated from equine infections at the University of Zurich between 1988 and 2014.

Material and Methods

Data collection

The database of the Institute of Veterinary Bacteriology and the Equine Hospital of the University of Zurich were retrospectively searched for bacterial isolates cultured from horses presented to the Equine Hospital of the University of Zurich. The information obtained included date of collection, organ system affected, bacterial strain isolated and number of pathogenic bacteria per sample.

Classification

Infections were classified into 12 groups based on the origin of the samples. Respiratory tract infections included pathogens isolated from the upper and lower airways, including pleural involvement. Implant infections (orthopedic), infections of incisions and injection sites (soft tissue) and infections after dental procedures (sinus) were grouped as post-procedural infections. Synovial infections, osteitis and hoof abscesses were combined to musculoskeletal infections. Soft tissue infections included external abscesses, wounds, cellulitis, infected hematomas and myositis. Isolates from the uterus, vagina, lochia, prepuce and samples from mastitis were summarized as reproductive infections. Urinary tract infections included upper and lower urinary tract infection and omphalitis. Dental infections were sampled from horses with clinical appearance of an alveolar periostitis, dental sinusitis or osteitis due to dental disease. Nervous system infections contained all isolates from cerebrospinal fluid. Bacterial isolates from undetermined sites and samples collected from lymph nodes, ear swabs, salivary glands, neoplasia and liver biopsies were classified as 'others'. Bacteria isolated from fecal samples were excluded. Isolates considered as non-pathogenic or contamination were excluded based on information on bacterial species, origin of sample, quantity of growth, the occurrence of mixed cultures and clinical findings. If opportunistic pathogens were isolated, they were only included if a role as causative pathogen seemed reasonable based on clinical findings. When mixed infections occurred in one sample, all isolates were included. When a bacterial strain was isolated more than once from the same site in a horse within six months, the isolate was only counted once in the study (persistent infection).

Period of time

The available data was divided into four phases: 1988–2002 (A1), 2003–2006 (A2), 2007–2010 (B1) and 2011–2014 (B2). The time spans were chosen to reflect major changes in antimicrobial treatment at the Equine Hospital. From 1988 to 2006 (A1 and A2), the main first-line antibiotic treatment choice was penicillin, normally combined with the aminoglycoside gentamicin. Due to a perceived increase of resistance against gentamicin, cefquinome was used more frequently as an agent between 2007 and 2014 (B1 and B2).

Culture methods

Depending on the isolation site, samples were streaked on standard agars (Columbia blood agar with sheep blood, Gassner agar, Columbia CNA agar, Chocolate agar with vitox, Oxoid AG, Pratteln, Switzerland) and incubated at 37°C for 24–48 hours under aerobic conditions. For culture of obligate anaerobes, samples were streaked on Columbia blood agar with sheep blood

Table 2: Bacterial isolates from horses in Zurich between 1988 and 2014.

Bacterial strains with less than 10 counts (<0.5%) grouped in 'others' included: *Bordetella bronchiseptica*, *Rhodococcus equi*, *Dermatophilus congolensis*, *Stenotrophomonas maltophilia*, *Nocardia* spp., *S. canis*, *S. uberis*, *Aeromonas* spp., *Mannheimia haemolytica*, *Neisseria* spp.
* Formerly *Arcanobacterium pyogenes*

| Bacterial isolate | n (%) |
|------------------------------------------------|-------------|
| <i>E. coli</i> | 299 (17.4%) |
| <i>S. equi</i> ssp. <i>zooepidemicus</i> | 295 (17.1%) |
| coagulase-positive staphylococci | 196 (11.4%) |
| coagulase-negative staphylococci | 133 (7.7%) |
| Anaerobes | 124 (7.2%) |
| <i>Enterococcus</i> spp. | 94 (5.5%) |
| <i>P. aeruginosa</i> | 87 (5.1%) |
| <i>Acinetobacter</i> spp. | 70 (4.1%) |
| <i>Pasteurella</i> spp. | 55 (3.2%) |
| <i>Enterobacter</i> spp. | 52 (3.0%) |
| <i>Klebsiella</i> spp. | 46 (2.7%) |
| <i>Proteus</i> spp. | 40 (2.3%) |
| <i>S. equi</i> ssp. <i>equi</i> | 35 (2.0%) |
| <i>Actinobacillus</i> spp. | 32 (1.9%) |
| <i>Corynebacterium</i> spp. | 26 (1.5%) |
| <i>Pantoea agglomerans</i> | 25 (1.5%) |
| <i>S. dysgalactiae</i> ssp. <i>equisimilis</i> | 24 (1.4%) |
| <i>Trueperella pyogenes</i> * | 23 (1.3%) |
| <i>Citrobacter</i> spp. | 14 (0.8%) |
| <i>Actinomyces</i> spp. | 11 (0.6%) |
| Total | 1723 (100%) |

and Schaedler KV agar (Oxoid AG, Pratteln, Switzerland) and incubated at 37°C under anaerobic conditions for 48–72 hours. The samples were investigated by standard bacteriological procedures using gram-staining, catalase test, oxidase test, oxidation-fermentation test, agglutination test to check coagulase activity, agglutination test to classify the Lancefield groups of *Streptococcus* spp. and growth performance on different agar (Markey, 2013). Identification to species level was done using the API® test system (1988–2011) or the VITEK® 2 compact system (2012–2014) (both bioMérieux, Marcy l'Etoile, France) using the appropriate API® strips or VITEK® cards according to the manufacturer's instructions.

Statistical analysis

To compare frequency distributions between different phases, the Chi-Square test or Fisher exact test was used. Statistical analysis was performed using commercial software (GraphPad Prism 6). The Bonferroni correction for multiple comparisons was applied and results were considered statistically significant if $p < 0.0125$.

A retrospective study of bacterial pathogens in an equine hospital (1988–2014)

J. N. van Spijk et al.

A retrospective study
of bacterial pathogens
in an equine hospital
(1988–2014)

J. N. van Spijk et al.




Figure 2: Changes in frequency of pathogenic bacterial strains over time isolated from horses between 1988 and 2014 (n=1,723).

* indicates a significant change over time, different letters show significant difference between two groups (p < 0.0125)

Results

The database yielded 1,723 bacterial pathogens isolated from 1,281 samples from horses between 1988 and 2014. During phase A1 (1988–2002) 258/1,723 (15%) isolates were obtained, during phase A2 (2003–2006) 481/1,723 (28%), during phase B1 (2007–2010) 445/1,723 (26%) and during phase B2 (2011–2014) 539/1,723 (31%). The majority of isolates was obtained from respiratory (397/1,723, 23%), post-procedural (386/1,723, 22%) and soft tissue infections (208/1,723, 12%). An overview of the origin of samples is shown in Table 1. In 940/1,281 (73%) samples a single bacterial species was identified, whereas in 346/1,281 (27%) samples mixed infections (2–6 isolates) occurred. In 206/346 (60%) mixed infections simultaneous infection with Gram-positive and Gram-negative bacteria was present. The occurrence of mixed infections varied between organ systems (Fig. 1).

The most commonly isolated bacteria were *Escherichia (E.) coli* (299/1,723, 17%), *Streptococcus (S.) equi* ssp. *zooepidemicus* (295/1,723, 17%) and coagulase-positive staphylococci (196/1,723, 11%). All isolated bacterial pathogens are listed in Table 2. Coagulase-positive staphylococci were grouped together, whereof 136/196 (69%) were

Staphylococcus aureus. Of *Acinetobacter* spp. 31/70 (44%) were further classified into *A. baumannii*. Aerobic bacteria dominated with 1,599/1,723 (93%) isolates: 859/1,599 (54%) were Gram-positive and 740/1,599 (46%) Gram-negative. Obligate anaerobic bacteria represented 124/1,723 (7%) isolates from 113 samples: 22/124 (18%) were Gram-positive, 67/124 (54%) were Gram-negative and 35/124 (28%) were not further classified. Mixed aerobic and anaerobe infections were present in 62/113 (55%) of samples. Most common obligate anaerobes were *Bacteroides* spp. (37/124, 30%), *Clostridium* spp. (18/124, 15%) and *Fusobacterium* spp. (18/124, 15%).

Statistical analysis was performed for bacterial strains isolated at least 10 times (>0.5%). Significant changes over time were identified for anaerobes (p<0.0001), coagulase-negative staphylococci (p<0.0001), *Enterococcus* spp. (p<0.0001), *Pseudomonas (P.) aeruginosa* (p<0.0001), *Acinetobacter* spp. (p=0.0079), *Pasteurella* spp. (p=0.0007), *Proteus* spp. (p=0.0065), *Actinobacillus* spp. (p=0.002), *Pantoea agglomerans* (p<0.0001), *S. dysgalactiae* ssp. *equisimilis* (p=0.0074), and *Trueperella* (formerly *Arcanobacterium*) *pyogenes* (p=0.0046). Both increases and decreases over time were observed (Fig. 2).

The distribution of the most common bacterial strains by organ system is shown in Table 3. Due to the small numbers of bacterial strains in several organ systems, longitudinal analysis of bacterial isolates was only performed for respiratory, post-procedural, soft tissue and musculoskeletal infections. Significant changes over time were seen in respiratory and post-procedural infections (Fig. 3).

Discussion

This study shows that *E. coli*, *S. equi* ssp. *zooepidemicus* and coagulase-positive staphylococci were the 3 most commonly isolated equine pathogens. In most organ systems Gram-negative as well as Gram-positive bacteria were common. These results agree with previous studies reporting *E. coli* and *S. equi* ssp. *zooepidemicus* as the most commonly isolated equine pathogens (Lavoie et al., 1991; Clark et al., 2008; Panchaud et al., 2010). Similar to an earlier study in another region of Switzerland (Panchaud et al., 2010), coagulase-positive staphylococci, including *S. aureus*, were particularly numerous, while *Actinobacillus* spp., an important pathogen in other countries, was uncommon (Lavoie et al., 1991; Clark et al., 2008). Differing bacterial prevalence between geographic regions has previously been described (Sellon and Long, 2013) and is likely due to differing environments, prophylactic and therapeutic measures, eradication strategies, biosecurity and antimicrobial use. This further highlights the need for prevalence studies in different regions.

S. equi ssp. *equi* is a common respiratory pathogen in Switzerland (Panchaud et al., 2010), showing that strangles is still widely prevalent in the Swiss horse population. *Rhodococcus equi*, an important respiratory pathogen of foals in geographically close regions (Venner et al., 2013), was only isolated from 5/397 (1%) samples. Anaerobes, often involved in cases of pleuropneumonia (Sweeney et al., 1991), were rarely isolated from respiratory infections, which corresponds with the fact that pleuropneumonia is an uncommon disease at our hospital. Frequent isolates from post-procedural infections in our and other studies (Clark et al., 2008; Taylor et al., 2010; Torfs et al., 2010) included *E. coli*, *Staphylococcus* spp., *Enterococcus* spp., *P. aeruginosa*, and *Acinetobacter* spp. These species are known for their ability to develop high rates of antimicrobial resistance, and multidrug-resistant isolates are reported to cause incisional and implant infections (Weese, 2009). Thus, treatment in post-procedural infections should always be guided by a susceptibility profile. In peritonitis cases, isolation of *E. coli* and anaerobes was common and in accordance with previous studies (Hirsh and Jang, 1987; Lavoie et al., 1991; Hawkins et al., 1993; Southwood and Russell,




Figure 3: Bacterial strains with significant changes over time cultured from equine (1) respiratory (n=397) and (2) post-procedural infections (n=386) in Zurich between 1988 and 2014.

Different letters show significant difference between two groups ($p < 0.0125$)

2007). Only a single case of unclassified *Actinobacillus* spp. occurred, while this pathogen is an important cause of equine peritonitis in Australia and America (Matthews et al., 2001; Southwood and Russell, 2007). In neonatal sepsis, *Enterobacteriaceae* and other Gram-

A retrospective study
of bacterial pathogens
in an equine hospital
(1988–2014)

J. N. van Spijk et al.

Table 3: Distribution of bacterial isolates (n=1,723) from horses in Zurich between 1988 and 2014 by organ system of origin. Pathogenic bacteria with an occurrence of >5% isolated from the organ system are shown.

| Origin of samples | Total number of isolates | Bacterial strain | n (%) |
|-------------------|--------------------------|------------------------------------------|----------|
| Respiratory | n=397 | <i>S. equi</i> ssp. <i>zooepidemicus</i> | 99 (25%) |
| | | <i>E. coli</i> | 58 (15%) |
| | | <i>S. equi</i> ssp. <i>equi</i> | 33 (8%) |
| | | <i>Pasteurella</i> spp. | 29 (7%) |
| Post-procedural | n=386 | <i>E. coli</i> | 83 (22%) |
| | | coagulase-positive staphylococci | 55 (14%) |
| | | coagulase-negative staphylococci | 51 (13%) |
| | | <i>Enterococcus</i> spp. | 30 (8%) |
| | | <i>P. aeruginosa</i> | 28 (7%) |
| | | <i>S. equi</i> ssp. <i>zooepidemicus</i> | 27 (7%) |
| | | <i>Acinetobacter</i> spp. | 26 (7%) |
| | | Anaerobes | 26 (7%) |
| Soft tissue | n=208 | <i>S. equi</i> ssp. <i>zooepidemicus</i> | 37 (18%) |
| | | coagulase-positive staphylococci | 36 (17%) |
| | | Anaerobes | 28 (13%) |
| | | <i>E. coli</i> | 21 (10%) |
| | | <i>Acinetobacter</i> spp. | 13 (6%) |
| | | coagulase-negative staphylococci | 12 (6%) |
| Musculoskeletal | n=207 | <i>S. equi</i> ssp. <i>zooepidemicus</i> | 35 (17%) |
| | | coagulase-positive staphylococci | 33 (16%) |
| | | <i>E. coli</i> | 30 (15%) |
| | | coagulase-negative staphylococci | 20 (10%) |
| | | Anaerobes | 14 (7%) |
| | | <i>P. aeruginosa</i> | 13 (6%) |
| | | <i>Proteus</i> spp. | 13 (6%) |
| | | <i>Enterococcus</i> spp. | 13 (6%) |
| Reproductive | n=99 | <i>S. equi</i> ssp. <i>zooepidemicus</i> | 35 (35%) |
| | | <i>E. coli</i> | 31 (31%) |
| | | <i>Enterobacter</i> spp. | 6 (6%) |
| Urinary | n=95 | <i>E. coli</i> | 30 (32%) |
| | | <i>Enterococcus</i> spp. | 16 (17%) |
| | | <i>S. equi</i> ssp. <i>zooepidemicus</i> | 12 (13%) |
| | | coagulase-negative staphylococci | 8 (8%) |
| | | coagulase-positive staphylococci | 6 (6%) |
| | | <i>Klebsiella</i> spp. | 6 (6%) |
| Skin | n=87 | coagulase-positive staphylococci | 29 (33%) |
| | | coagulase-negative staphylococci | 10 (11%) |
| | | <i>P. aeruginosa</i> | 10 (11%) |
| | | <i>S. equi</i> ssp. <i>zooepidemicus</i> | 8 (9%) |
| | | <i>Dermatophilus congolensis</i> | 5 (6%) |
| | | <i>Pantoea agglomerans</i> | 5 (6%) |
| Dental | n=66 | Anaerobes | 18 (27%) |
| | | <i>E. coli</i> | 13 (20%) |
| | | <i>S. equi</i> ssp. <i>zooepidemicus</i> | 10 (15%) |
| | | <i>P. aeruginosa</i> | 5 (8%) |
| | | <i>Enterococcus</i> spp. | 5 (8%) |
| Peritoneal | n=43 | <i>E. coli</i> | 13 (30%) |
| | | Anaerobes | 6 (14%) |
| | | <i>S. equi</i> ssp. <i>zooepidemicus</i> | 5 (12%) |
| | | <i>Enterococcus</i> spp. | 5 (13%) |
| | | coagulase-positive staphylococci | 3 (7%) |
| Sepsis | n=27 | <i>E. coli</i> | 8 (30%) |
| | | coagulase-negative staphylococci | 5 (19%) |
| | | <i>Pasteurella</i> spp. | 4 (15%) |
| | | <i>Actinobacillus</i> spp. | 2 (7%) |
| | | <i>Klebsiella</i> spp. | 2 (7%) |

| Origin of samples | Total number of isolates | Bacterial strain | n (%) |
|-------------------|--------------------------|------------------------------------------|----------|
| Nervous system | n=3 | <i>Acinetobacter</i> spp. | 1 (33%) |
| | | <i>E. coli</i> | 1 (33%) |
| | | <i>Enterococcus</i> spp. | 1 (33%) |
| Others | n=105 | <i>S. equi</i> ssp. <i>zooepidemicus</i> | 27 (26%) |
| | | <i>Actinobacillus</i> spp. | 11 (10%) |
| | | coagulase-positive staphylococci | 9 (9%) |
| | | <i>E. coli</i> | 9 (9%) |
| | | <i>Pasteurella</i> spp. | 8 (8%) |
| | | <i>P. aeruginosa</i> | 7 (7%) |
| | | Anaerobes | 6 (6%) |

A retrospective study of bacterial pathogens in an equine hospital (1988–2014)

J. N. van Spijk et al.

negatives are the most common isolates (Lavoie et al., 1991; Clark et al., 2008; Theelen et al., 2014). However, an increase in Gram-positive isolates, especially *Enterococcus* spp., has been reported recently (Theelen et al., 2014). Sepsis cases in our study showed a similar spectrum, but the case numbers were too low for longitudinal analysis. Nervous system infection in foals is usually a sequel to sepsis (Tyler et al., 1993; Pusterla et al., 2007; Hepworth et al., 2014). In this study *E. coli* was isolated from the cerebrospinal fluid of a foal with sepsis and a corresponding positive blood culture. In contrast, little is known about bacterial agents causing meningoencephalitis in adult horses. Only 2 cases of adult horses with clinical signs of meningoencephalitis and a positive culture result for *Acinetobacter* spp. and *Enterococcus* spp., respectively, occurred over the study period. Prediction of the etiologic pathogen is therefore difficult in these cases, highlighting the need for bacteriologic culture.

As reported before (Lavoie et al., 1991), most samples (935/1,281, 73%) yielded pure cultures of one sole bacterial pathogen. Mixed infections were particularly common in organs colonized with a large number of commensal bacteria like the skin or the oral cavity.

While the occurrence of main pathogens remained stable, some bacterial strains emerged and others were less commonly isolated over the last years, displaying a dynamic interaction between horse population and pathogens adapting to their environment. Antimicrobial therapy has an impact on bacterial prevalence, as it allows bacteria to grow which are intrinsically resistant or able to develop resistance to the used antimicrobial drug (Morley et al., 2005; Dunowska et al., 2006). Some antimicrobial classes have a greater potential to promote selection of resistant isolates than others, as shown for cephalosporins in equine fecal *E. coli* (Dunowska et al., 2006). The change of first-line antibiotics from penicillin/gentamicin to cefquinome in 2007 could have influenced the bacterial spectrum. The increased isolation rates of *Enterococcus* spp., *Acinetobacter* spp., and coagulase-negative staphylococci possibly reflect their ability to

adapt quickly and develop resistance to several antimicrobial drugs. In addition, *Enterococcus* spp. are intrinsically resistant to all cephalosporins (Weese, 2009), which could further explain their increasing occurrence during the last years. Multidrug-resistant enterococci (primarily *E. faecalis* and *E. faecium*), *Acinetobacter* spp. (especially *A. baumannii*), and coagulase-negative staphylococci (especially methicillin-resistant staphylococci) are increasingly important nosocomial pathogens in humans (Weese, 2009; Becker et al., 2014). Little is known about such infections in horses, but their increasing occurrence seen in hospitalized horses in this study is alarming.

One limitation of this study is its retrospective nature. Changes in laboratory techniques, clinical diagnostics and knowledge of causative organisms occurred over the time investigated. Awareness of important isolates and new possibilities of classifying them may increase the number of positive results without a real increase in their prevalence. Furthermore, a true prevalence for each bacterial pathogen could not be calculated, because some horses with clinical signs of an infection might not have had a culture taken and samples revealing no growth were excluded. Care should be taken when extrapolating the results to private practice, as all our samples were taken from hospitalized horses and probably many of them were already pretreated with antibiotics.

Conclusion

Gram-positive as well as Gram-negative bacteria were common isolates in almost all organ systems. Furthermore, in dental, peritoneal and soft tissue infections anaerobic bacteria should be considered. In conclusion, initial broad-spectrum antimicrobial therapy is generally necessary until bacteriological culture and susceptibility testing results become available. However, clinical diagnosis may allow targeted antimicrobial therapy in some cases (e.g. strangles). Longitudinal changes reflect a dynamic interaction between pathogens, environment and host. Antimicrobial therapy does influence

A retrospective study
of bacterial pathogens
in an equine hospital
(1988–2014)

J. N. van Spijk et al.

bacterial prevalence as it selects for isolates resistant to the used antibiotic. Monitoring of bacterial pathogens in horses should be continued and susceptibility testing data of reported pathogens analyzed.

Acknowledgements

The study was supported by a Forschungskredit of the University of Zurich, grant no. [FK-14-052].

Etude rétrospective des bactéries pathogènes sur des chevaux hospitalisés (1988–2014)

Une connaissance des bactéries pathogènes présentes localement est importante pour pouvoir mettre en œuvre dès le début un traitement antibiotique ciblé. Le but de cette étude était de décrire les agents infectieux les plus fréquents et leur évolution dans le temps chez les chevaux hospitalisés à l'hôpital vétérinaire de Zurich entre 1988 et 2014. Tous les résultats positifs de cultures ont été analysés rétrospectivement. Les isolats ont été classés sur la base de leur localisation, du diagnostic et de l'année de leur isolation. Diverses périodes ont été comparées au moyen d'un test de chi carré. *Escherichia coli* (299/1'723, 17%), *Streptococcus equi* ssp. *zooepidemicus* (295/1'723, 17%) et les staphylocoques coagulase positive (196/1'723, 11%) ont été dans l'ensemble les bactéries mises en évidence le plus souvent. Des bactéries anaérobies strictes ont été trouvées principalement dans les infections dentaires (18/66, 27%), les péritonites (6/43, 14%) et les infections des tissus mous (28/208, 13%). Des cultures mixtes étaient particulièrement fréquentes dans les infections cutanées et dentaires (24/55, 44% et 17/43, 40%). Les principaux germes isolés dans la plupart des systèmes d'organes étaient aussi bien Gram positifs que Gram négatifs. Pour cette raison, il conviendrait, jusqu'à ce que les résultats des cultures et des antibiogrammes soient connus, de choisir un médicament à large spectre d'action.

Studio retrospettivo dei batteri patogeni nei cavalli ospedalizzati (1988–2014)

Le conoscenze sui batteri patogeni locali sono importanti al fine di avviare una terapia antibiotica mirata. Lo scopo dello studio era di descrivere gli agenti infettivi più comuni e le loro variazioni nel tempo nei cavalli presso il Tierspital Zürich (Ospedale Veterinario di Zurigo) tra il 1988 e il 2014. Tutti i risultati delle colture positivi sono stati analizzati retrospettivamente. Gli isolati sono stati classificati per la loro localizzazione, la diagnosi e l'anno di isolamento. Diversi periodi di tempo sono stati confrontati utilizzando il test chi quadrato. *Escherichia coli* (299/1'723, 17%), *Streptococcus equi* ssp. *zooepidemicus* (295/1'723, 17%) e stafilococchi coagulasi-positivi (196/1'723, 11%) sono stati i batteri più frequentemente rilevati globalmente. Batteri anaerobi obbligati sono stati isolati principalmente nelle infezioni ai denti (18/66, 27%), nella peritonite (6/43, 14%) e nelle infezioni dei tessuti molli (28/208, 13%). Colture miste erano particolarmente comuni nelle infezioni della pelle e dei denti (24/55, 44% e 17/43, 40%). In molti sistemi di organi, i batteri patogeni isolati più di frequente erano i gram-positivi e i gram-negativi. Pertanto, la scelta dell'antibiotico dovrebbe portare su un farmaco ad ampio spettro in attesa dei risultati delle colture e degli antibiogrammi.

References

- Becker, K., Heilmann, C., Peters, G.: Coagulase-negative staphylococci. Clin. Microbiol. Rev. 2014, 27: 870–926.
- Clark, C., Greenwood, S., Boison, J. O., Chirino-Trejo, M., Dowling, P. M.: Bacterial isolates from equine infections in western Canada (1998–2003). Can. Vet. J. 2008, 49: 153–160.
- Dunowska, M., Morley, P. S., Traub-Dargatz, J. L., Hyatt, D. R., Dargatz, D. A.: Impact of hospitalization and antimicrobial drug administration on antimicrobial susceptibility patterns of commensal *Escherichia coli* isolated from the feces of horses. J. Am. Vet. Med. Assoc. 2006, 228: 1909–1917.
- Hawkins, J. F., Bowman, K. F., Roberts, M. C., Cowen, P.: Peritonitis in horses: 67 cases (1985–1990). J. Am. Vet. Med. Assoc. 1993, 203: 284–288.
- Hepworth, K. L., Wong, D. M., Sponseller, B. A., Alcott, C. J., Sponseller, B. T., Ben-Shlomo, G., Whitley, R. D.: Survival of an adult Quarter Horse gelding following bacterial meningitis caused by *Escherichia coli*. Equine Veterinary Education. 2014, 26: 507–512.
- Hirsh, D. C., Jang, S. S.: Antimicrobial susceptibility of bacterial pathogens from horses. Vet. Clin. North. Am. Equine Pract. 1987, 3: 181–190.
- Lavoie, J. P., Couture, L., Higgins, R., Laverty, S.: Aerobic bacterial isolates in horses in a university hospital, 1986–1988. Can. Vet. J. 1991, 32: 292–294.
- Markey, B. K.: Clinical veterinary microbiology, 2nd Edition. Elsevier, Edinburgh.
- Matthews, S., Dart, A. J., Dowling, B. A., Hodgson, J. L., Hodgson, D. R.: Peritonitis associated with *Actinobacillus equuli* in horses: 51 cases. Aust. Vet. J. 2001, 79: 536–539.
- Morley, P. S., Apley, M. D., Besser, T. E., Burney, D. P., Fedorka-Cray, P. J., Papich, M. G., Traub-Dargatz, J. L., Weese, J. S.: Antimicrobial Drug Use in Veterinary Medicine. Journal of Veterinary Internal Medicine. 2005, 19: 617–629.
- Panchaud, Y., Gerber, V., Rossano, A., Perreten, V.: Bacterial infections in horses: a retrospective study at the University Equine Clinic of Bern. Schweiz. Arch. Tierheilk. 2010, 152: 176–182.
- Pusterla, N., Luff, J. A., Myers, C. J., Vernau, W., Affolter, V. K.: Disseminated Intravascular Coagulation in a Horse with *Streptococcus equi* subspecies zooepidemicus Meningoencephalitis and Interstitial Pneumonia. Journal of Veterinary Internal Medicine. 2007, 21: 344–347.
- Racklyeft, D. J., Love, D. N.: Bacterial infection of the lower respiratory tract in 34 horses. Aust. Vet. J. 2000, 78: 549–559.
- Sellon, D. C., Long, M. T.: Equine infectious diseases, 2nd Edition. Saunders/Elsevier, St. Louis, Missouri.
- Southwood, L. L., Russell, G.: The use of clinical findings in the identification of equine peritonitis cases that respond favorably to medical therapy. Journal of Veterinary Emergency and Critical Care. 2007, 17: 382–390.
- Sweeney, C. R., Holcombe, S. J., Barningham, S. C., Beech, J.: Aerobic and anaerobic bacterial isolates from horses with pneumonia or pleuropneumonia and antimicrobial susceptibility patterns of the aerobes. J. Am. Vet. Med. Assoc. 1991, 198: 839–842.
- Taylor, A. H., Mair, T. S., Smith, L. J., Perkins, J. D.: Bacterial culture of septic synovial structures of horses: does a positive bacterial culture influence prognosis? Equine Vet. J. 2010, 42: 213–218.
- Theelen, M. J., Wilson, W. D., Edman, J. M., Magdesian, K. G., Kass, P. H.: Temporal trends in prevalence of bacteria isolated from foals with sepsis: 1979–2010. Equine Vet. J. 2014, 46: 169–173.
- Torfs, S., Levet, T., Delesalle, C., Dewulf, J., Vlamincx, L., Pille, F., Lefere, L., Martens, A.: Risk factors for incisional complications after exploratory celiotomy in horses: do skin staples increase the risk? Vet. Surg. 2010, 39: 616–620.
- Traub-Dargatz, J. L., Dargatz, D. A.: Antibacterial drug resistance and equine practice. Equine Veterinary Education. 2009, 21: 49–56.
- Tyler, C. M., Davis, R. E., Begg, A. P., Hutchins, D. R., Hodgson, D. R.: A survey of neurological diseases in horses. Aust. Vet. J. 1993, 70: 445–449.
- Venner, M., Astheimer, K., Lämmer, M., Giguère, S.: Efficacy of Mass Antimicrobial Treatment of Foals with Subclinical Pulmonary Abscesses Associated with *Rhodococcus equi*. Journal of Veterinary Internal Medicine. 2013, 27: 171–176.
- Weese, J. S.: Antimicrobial therapy for multidrug resistant pathogens. Equine Veterinary Education. 2009, 21: 328–334.
- Wilson, W. D.: Rational Selection for Antimicrobials for Use in Horses. Proc. 47th Annu. Conv. Am. Assoc. Equine Pract. 2001, 47: 75–93.

A retrospective study of bacterial pathogens in an equine hospital (1988–2014)

J. N. van Spijk et al.

Corresponding author

Angelika Schoster
Dr. med. vet., DVSc, PhD, DACVIM/DECEIM
Oberärztin
Universität Zürich
Vetsuisse-Fakultät
Klinik für Pferdemedizin
Winterthurerstrasse 260
8057 Zürich
Schweiz
Tel. 044 635 83 57
E-Mail: aschoster@vetclinics.uzh.ch